



Table 2.3 Atomic Structures of the Most Abundant Elements in the Body					
Element	Symbol	Atomic number (# of p)	Mass number $(# of p + n)$	Atomic weight	Electrons in valence shell
Calcium	Ca	20	40	40.078	2
Carbon	С	6	12	12.011	4
Chlorine	Cl	17	35	35.453	7
Hydrogen	Н	1	1	1.008	1
lodine	I	53	127	126.905	7
Iron	Fe	26	56	55.847	2
Magnesium	Mg	12	24	24.305	2
Nitrogen	N	7	14	14.007	5
Oxygen	0	8	16	15.999	6
Phosphorus	Р	15	31	30.974	5
Sodium	Na	11	23	22.989	1
Sulfur	ς	16	32	32.064	6

© 2018 Pearson Education, Inc.




(a) Chemically inert elements

Outermost energy level (valence shell) complete



Helium (He) (2p+; 2n0; 2e-)



Neon (Ne) (10p+; 10n0; 10e-)

© 2018 Pearson Education, Inc.


7

(b) Chemically reactive elements


Outermost energy level (valence shell) incomplete




Hydrogen (H) (1p+; 0n0; 1e-)



Oxygen (O) (8p+; 8n0; 8e-)




Carbon (C) (6p+; 6n0; 6e-)




Sodium (Na) (11p+; 12n⁰; 11e-)

© 2018 Pearson Education, Inc.

8








Resulting molecules

Molecule of oxygen gas (O₂)



(b) Formation of a double covalent bond

Oxygen atom


© 2018 Pearson Education, Inc.

Oxygen atom

11


Reacting atoms


Resulting molecules





Hydrogen atoms Carbon atom Molecule of methane gas (CH₄)


(c) Formation of four single covalent bonds

























(b) Double sugar (disaccharide)

© 2018 Pearson Education, Inc.

2




(c) Dehydration synthesis and hydrolysis of a molecule of sucrose




(d) Starch (polysaccharide)

© 2018 Pearson Education, Inc.

23




(a) Saturated fat. At room temperature, the molecules of a saturated fat such as this butter are packed closely together, forming a solid.




(b) Unsaturated fat. At room temperature, the molecules of an unsaturated fat such as this olive oil cannot pack together closely enough to solidify because of the kinks in some of their fatty acid chains.


© 2018 Pearson Education, Inc.

(b) Typical structure of a phospholipid molecule (phosphatidylcholine). Two fatty acid chains and a phosphorous-containing group are attached to a glycerol backbone.



(c) Cholesterol. Simplified structure of cholesterol, a steroid, formed by four interlocking carbon rings.






(a) **Primary structure.** A protein's primary structure is the unique sequence of amino acids in the polypeptide chain.

29


© 2018 Pearson Education, Inc.



© 2018 Pearson Education, Inc.

by hydrogen bonds, represented by dashed lines in this figure.


alpha-helix and beta-pleated sheet. Secondary structure is reinforced



Protein (if > 50 amino acids) or polypeptide (if < 50 amino acids)

(c) Tertiary structure. The overall three-dimensional shape of the polypeptide or protein is called tertiary structure. It is reinforced by chemical bonds between the R-groups of amino acids in different regions of the protein chain.

© 2018 Pearson Education, Inc.




Complex protein with four polypeptide subunits, each with tertiary structure

(d) Quaternary structure. Some proteins consist of two or more polypeptide chains. For example, four polypeptides construct the protein hemoglobin, a blood protein.





(a) Triple helix of collagen (a fibrous or structural protein).


© 2018 Pearson Education, Inc.





(b) Hemoglobin molecule composed of the protein globin and attached heme groups. (Globin is a globular, or functional, protein.)













(a) Chemical work. ATP provides the energy needed to drive energy-absorbing chemical reactions.



(b) Transport work. ATP drives the transport of certain solutes (amino acids, for example) across cell membranes.



(c) Mechanical work. ATP activates contractile proteins in muscle cells so that the cells can shorten and perform mechanical work.