


1


Name	Molecular Formula	Structural Formula	Ball-and-Stick Model	Space-Filling Model
(a) Methane	CH ₄	$\begin{array}{c} \text{H} \\ \\ \text{H}-\text{C}-\text{H} \\ \\ \text{H} \end{array}$		
(b) Ethane	C ₂ H ₆	$\begin{array}{cc} \text{H} & \text{H} \\ & \\ \text{H}-\text{C} & -\text{C}-\text{H} \\ & \\ \text{H} & \text{H} \end{array}$		
(c) Ethene (ethylene)	C ₂ H ₄	$\begin{array}{cc} \text{H} & \text{H} \\ & \backslash \text{ } / \\ & \text{C}=\text{C} \\ & / \text{ } \backslash \\ \text{H} & \text{H} \end{array}$		

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.


2




3






4



5



6




Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.

7

Drug	Condition	Effective Enantiomer	Ineffective Enantiomer
Ibuprofen	Pain; inflammation	 S-Ibuprofen	 R-Ibuprofen
Albuterol	Asthma	 R-Albuterol	 S-Albuterol


Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.

8



Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.

9



Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.

10

CHEMICAL GROUP	Hydroxyl	Carbonyl	Carboxyl
STRUCTURE			
NAME OF COMPOUND	Alcohols (their specific names usually end in -ol)	Ketones if the carbonyl group is within a carbon skeleton. Aldehydes if the carbonyl group is at the end of the carbon skeleton.	Carboxylic acids, or organic acids
EXAMPLE	 Ethanol, the alcohol present in alcoholic beverages	 Acetone, the simplest ketone Propanal, an aldehyde	 Acetic acid, which gives vinegar its sour taste
FUNCTIONAL PROPERTIES	<ul style="list-style-type: none"> Is polar as a result of the electronegative oxygen atom. Can form hydrogen bonds with water molecules, helping dissolve organic compounds such as sugars. 	<ul style="list-style-type: none"> A ketone and an aldehyde may be structural isomers with different properties, as is the case for acetone and propanal. These two groups are also found in sugars, giving rise to two major groups of sugars: aldohexoses (containing an aldehyde) and ketohexoses (containing a ketone). 	<ul style="list-style-type: none"> Has acidic properties because the covalent bond between oxygen and hydrogen is so polar, for example, $\text{H}-\text{O}-\text{C}(=\text{O})-\text{R} \rightleftharpoons \text{H}^+ + \text{O}^--\text{C}(=\text{O})-\text{R}$. Found in cells in the ionized form with a charge of -1 and called a carboxylate ion (note, specifically, the acetate ion).


Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.

11

CHEMICAL GROUP	Amino	Sulphydryl	Phosphate	Methyl
STRUCTURE				
NAME OF COMPOUND	Amines	Thiols	Organic phosphates	Methylated compounds
EXAMPLE	 Glycine Because it also has a carboxyl group, glycine is both an amino and a carboxylic acid. Compounds with both groups are called amino acids.	 Cysteine Cysteine is an important sulfur-containing amino acid.	 Glycerol phosphate In addition to taking part in many important chemical reactions in cells, glycerol phosphate provides the backbone for phospholipids, the most prevalent molecules in cell membranes.	 5-Methyl cytosine 5-Methyl cytosine is a component of DNA that has been modified by addition of the methyl group.
FUNCTIONAL PROPERTIES	<ul style="list-style-type: none"> Acts as a base; can pick up an H^+ from the surrounding solution (note, living organisms). Cross-linking of cysteines in hair proteins maintains the curliness or straightness of hair. Straight hair can be "permanently" curled by adding a sulfur carrier, then breaking and reforming the cross-linking bonds. 	<ul style="list-style-type: none"> Two sulphydryl groups can react, forming a covalent bond. This "cross-linking" helps stabilize protein structure. 	<ul style="list-style-type: none"> Contributes negative charge to the molecule of which it is a part (2- when at the end of a molecule, 1- when located internally in a chain of phosphates). Has the potential to react with water, releasing energy. 	<ul style="list-style-type: none"> Addition of a methyl group to DNA, as by molecules bound to DNA, affects expression of genes. Arrangement of methyl groups in male and female sex hormones affects their shape and function.

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.

12



Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.