



Figure 5-1



Copyright © 2010 Pearson Education, Inc.

1


Figure 5-2



Copyright © 2010 Pearson Education, Inc.

2


Figure 5-3



Copyright © 2010 Pearson Education, Inc.

3


Figure 5-4



Copyright © 2010 Pearson Education, Inc.

4


Figure 5-5



Copyright © 2010 Pearson Education, Inc.

5

Figure 5-6, overview



Copyright © 2010 Pearson Education, Inc.

6

Table 5-1, part 1

TABLE 5-1 Rules for Diffusion of Uncharged Molecules**General Properties of Diffusion**

1. Diffusion uses the kinetic energy of molecular movement and does not require an outside energy source.
2. Molecules diffuse from an area of higher concentration to an area of lower concentration.
3. Diffusion continues until concentrations come to equilibrium. Molecular movement continues, however, after equilibrium has been reached.
4. Diffusion is faster
 - along higher concentration gradients.
 - over shorter distances.
 - at higher temperatures.
 - for smaller molecules.
5. Diffusion can take place in an open system or across a partition that separates two systems.

Copyright © 2010 Pearson Education, Inc.

7

Table 5-1, part 2


TABLE 5-1 Rules for Diffusion of Uncharged Molecules (*continued*)**Simple Diffusion Across a Membrane**

6. The rate of diffusion through a membrane is faster if
 - the membrane's surface area is larger.
 - the membrane is thinner.
 - the concentration gradient is larger.
 - the membrane is more permeable to the molecule.
7. Membrane permeability to a molecule depends on
 - the molecule's lipid solubility.
 - the molecule's size.
 - the lipid composition of the membrane.

Copyright © 2010 Pearson Education, Inc.

8


Figure 5-7, overview



Copyright © 2010 Pearson Education, Inc.

9


Figure 5-8



Copyright © 2010 Pearson Education, Inc.

10


Figure 5-9, overview



Copyright © 2010 Pearson Education, Inc.

11


Figure 5-11



Copyright © 2010 Pearson Education, Inc.

12

Figure 5-12



Copyright © 2010 Pearson Education, Inc.

13

Table 5-2

TABLE 5-2


Primary Active Transporters

NAMES	TYPE OF TRANSPORT
Na^+/K^+ -ATPase or sodium-potassium pump	Antiport
Ca^{2+} -ATPase	Uniport
H^+ -ATPase or proton pump	Uniport
H^+/K^+ -ATPase	Antiport

Copyright © 2010 Pearson Education, Inc.

14


Figure 5-13



Copyright © 2010 Pearson Education, Inc.

15


Figure 5-14, overview



Copyright © 2010 Pearson Education, Inc.

16


Figure 5-15, overview



Copyright © 2010 Pearson Education, Inc.

17


Figure 5-16, overview



Copyright © 2010 Pearson Education, Inc.

18


Figure 5-18



Copyright © 2010 Pearson Education, Inc.

19


Figure 5-17



Copyright © 2010 Pearson Education, Inc.

20


Figure 5-19



Copyright © 2010 Pearson Education, Inc.

21


Figure 5-20, overview



Copyright © 2010 Pearson Education, Inc.

22


Figure 5-21, overview



Copyright © 2010 Pearson Education, Inc.

23


Figure 5-24



Copyright © 2010 Pearson Education, Inc.

24


Figure 5-25



Copyright © 2010 Pearson Education, Inc.

25


Figure 5-26, overview



Copyright © 2010 Pearson Education, Inc.

26


Figure 5-27



Copyright © 2010 Pearson Education, Inc.

27

Figure 5-28



Copyright © 2010 Pearson Education, Inc.

28

Table 5-5

TABLE 5-5

Comparing Osmolarities

SOLUTION A = 1 OsM GLUCOSE	SOLUTION B = 2 OsM GLUCOSE	SOLUTION C = 1 OsM NaCl
A is hypotonic to B	B is hyperosmotic to A	C is isotonic to A
A is isotonic to C	B is hyperosmotic to C	C is hypotonic to B

Copyright © 2010 Pearson Education, Inc.

29

Table 5-6

TABLE 5-6


Tonicity of Solutions

SOLUTION	CELL BEHAVIOR WHEN PLACED IN THE SOLUTION	DESCRIPTION OF THE SOLUTION RELATIVE TO THE CELL
A	Cell swells	Solution A is hypotonic
B	Cell doesn't change size	Solution B is isotonic
C	Cell shrinks	Solution C is hypertonic

Copyright © 2010 Pearson Education, Inc.

30

Figure 5-29




Copyright © 2010 Pearson Education, Inc.

31

Figure 5-29a

(a) Cell and solution are electrically and chemically at equilibrium.




Copyright © 2010 Pearson Education, Inc.

32

Figure 5-29b


(b) Cell and solution in chemical and electrical disequilibrium.



Copyright © 2010 Pearson Education, Inc.

33

Figure 5-29c




(c) On an absolute charge scale, the extracellular fluid (ECF) would be at +1 and the intracellular fluid (ICF) at -1.

Copyright © 2010 Pearson Education, Inc.

34


Figure 5-30



Copyright © 2010 Pearson Education, Inc.

35

Figure 5-33



Copyright © 2010 Pearson Education, Inc.

36

Figure 5-34




Figure 5-35

